



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

seeds, and there have been already matured or on the road to maturity 372 seed vessels, with 11,160 seeds. How many of these would get through the long chapter of accidents and produce flowering plants next year? I venture to say not a hundred—possibly not ten. What chance has an “occasional cross” to benefit the race in a scheme like that proposed?

And then we find that those which get more than an “occasional cross” do not get along any better for it. Take Mr. Robertson’s illustrations again. *Gaura biennis* I believe to be more dependent on insect aid than he himself has discovered, though none of those he names have any hand whatever in it, while its close ally *Gaura parviflora* is just as absolute a self fertilizer. And if *Oenothera fruticosa* is so arranged that self-pollination is impossible—a fact of which I am by no means sure—how about its neighbor *Oenothera biennis*, which is one of the closest self-fertilizers in the whole family, and yet has made its way not only all over the American continent, but has invaded the old world as well!

I repeat, where does the physiological advantage of the “occasional cross” come in?—THOMAS MEEHAN, *Germantown, Philadelphia*.

*Sullivantia Hapemaii*.—In the November GAZETTE (p. 348), owing to undue haste in printing, this species appeared as a *Heuchera*. The oversight was unfortunate, but it is to be hoped that the correction can overtake the blunder. It is a matter of some interest to discover in our flora a third species of *Sullivantia*, and that, too, with range intermediate between that of the other two. *S. Ohionis* of the north central states (Ohio to Iowa and Minnesota) has always been considered a rare and interesting plant; and *S. Oregana* of the Willamette and Columbia Rivers still more so. This third species, from the Big Horn Mountains of Wyoming, well preserves the generic appearance, and would be recognized at a glance by those familiar with the other species, although much more closely related to its eastern congener, a thing to be expected. All three species affect the same situation; all being found growing in the crevices of dripping cliffs. In addition to the description in the November GAZETTE it may be added that the calyx-lobes are 3-nerved and bright green; the petals are obovate, entire, and brown at base; and the pod is broad and depressed at the partition. The species somewhat resembles *S. Ohionis*, but its lower habit, smaller and deeply cut leaves (the lobes acutely dentate), green and 3-nerved calyx-lobes, much broader obovate (not oblanceolate acutish) petals, and its broad and depressed pod make it very distinct.—JOHN M. COULTER, *Bloomington, Ind.*